

Dr. Duke's Phytochemical and Ethnobotanical Databases

List of Plants for TRICYCLENE

Plant	Part	Low PPM	High PPM	StdDev	Reference
<i>Abies alba</i>	Leaf	45.0	63.0	0.5496290114333532	--
<i>Achillea millefolium</i>	Leaf	0.6	27.0	-0.8389074385035388	Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
<i>Artemisia salsoloides</i>	Shoot		1250.0	3.997049625995964	V. Kaul, P. Weyerstahl, H. Wahlberg, H. Marschall, (1992); Volatile constituents of the essential oil and the absolute of <i>Artemisia salsoloides</i> Willd. from Ladakh, Flavour and Fragrance journal, Vol.7, 299-305.
<i>Carum carvi</i>	Fruit				--
<i>Carum carvi</i>	Seed				Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
<i>Crocus sativus</i>	Silk Stigma Style				--
<i>Cymbopogon nardus</i>	Plant	45.0	180.0		Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
<i>Ferula gummosa</i>	Plant				--
<i>Juniperus communis</i>	Fruit				--
<i>Lavandula x intermedia</i>	Plant				Lawrence, B.M., Essential Oils 1976-1977, Essential Oils 1978, Essential Oils 1979-1980.
<i>Ocimum gratissimum</i>	Shoot		0.1	-0.2826219084407503	Vostrowsky, O., Garbe, W., Bestmann, H.J. and Maia, J.G.S. 1990. Essential Oil of Alfavaca, <i>Ocimum gratissimum</i> , from Brazilian Amazon. Zeitschr. Naturforschung 45(C): 1073-6.
<i>Ocimum basilicum</i>	Essential Oil				--
<i>Origanum sipyleum</i>	Shoot				Baser, K.H.C., Ozek, T., Kurkcuglu, M. and Tumen, G. 1992. Composition of the Essential Oil of <i>Origanum sipyleum</i> of Turkish Origin. J. Ess. Oil Res. 4: 139-142.
<i>Origanum sipyleum</i>	Shoot		30.0	-0.1802439751023571	Baser, K.H.C., Ozek, T., Kurkcuglu, M. and Tumen, G. 1992. Composition of the Essential Oil of <i>Origanum sipyleum</i> of Turkish Origin. J. Ess. Oil Res. 4: 139-142.

Plant	Part	Low PPM	High PPM	StdDev	Reference
Origanum onites	Shoot		15.0	-0.2316041423289758	Biondi, D., Cianci, P., Geraci, C. and Ruberto, G. 1993. Antimicrobial Activity and Chemical Composition of Essential Oils from Sicilian Aromatic Plants. <i>Flav. & Frag. J.</i> 8: 331-7.
Origanum syriacum	Shoot		3.0	-0.27269227611027075	Fleisher, A. & Fleisher, Z. 1991. Chemical Composition of <i>Origanum syriacum</i> L. Essential Oil. <i>J. Ess. Oil Res.</i> 3: 121-123.
Origanum sipyleum	Shoot		30.0	-0.1802439751023571	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of <i>Origanum sipyleum</i> of Turkish Origin. <i>J. Ess. Oil Res.</i> 4: 139-142.
Origanum syriacum	Shoot		0.0	-0.2829643095555945	Fleisher, A. & Fleisher, Z. 1991. Chemical Composition of <i>Origanum syriacum</i> L. Essential Oil. <i>J. Ess. Oil Res.</i> 3: 121-123.
Origanum sipyleum	Shoot		20.0	-0.21448408658676957	Baser, K.H.C., Ozek, T., Kurkcuoglu, M. and Tumen, G. 1992. Composition of the Essential Oil of <i>Origanum sipyleum</i> of Turkish Origin. <i>J. Ess. Oil Res.</i> 4: 139-142.
Origanum minutiflorum	Shoot	5.0	7.0	-0.25899623151650575	Baser, K.H.C., Tumen, G., Sezik, E. 1991. The Essential Oil of <i>Origanum minutiflorum</i> O. Schwarz and P.H. Davis. <i>J. Ess. Oil Res.</i> 3: 445-446.
Origanum syriacum	Shoot		0.0	-0.2829643095555945	Fleisher, A. & Fleisher, Z. 1991. Chemical Composition of <i>Origanum syriacum</i> L. Essential Oil. <i>J. Ess. Oil Res.</i> 3: 121-123.
Picea mariana	Twig		46.0		--
Pinus sylvestris	Resin, Exudate, Sap				--
Pinus sylvestris	Leaf Essent. Oil		4400.0	1.0	--
Pinus sylvestris	Essential Oil				--
Rosmarinus officinalis	Leaf Essent. Oil				--
Rosmarinus officinalis	Plant				Duke, J. A. Writeups or information summaries on approximately 2,000 economic plants, USDA, ARS, Beltsville, MD 20705.
Salvia officinalis	Leaf	15.0	84.0	1.3596086072298736	Duke, J. A. Writeups or information summaries on approximately 2,000 economic plants, USDA, ARS, Beltsville, MD 20705.

Plant	Part	Low PPM	High PPM	StdDev	Reference
<i>Salvia officinalis</i>	Leaf Essent. Oil		3000.0	-1.0	--
<i>Salvia canariensis</i>	Leaf		21.0	-1.0703301801596874	Casnigueral,S., Iglesias,J., Vila,R., Virgili,A. and Ibanez,C.1994. The Essential Oil from Leaves of <i>Salvia canariensis</i> L. Flav. & Fragr. J. 9:201-204. S. Canigueral, Facultat de Farmacia, Universitat de Barcelona, Ave.Diagonal 643,E-08028, Barcelone Spain
<i>Thymus riatarum</i>	Shoot		0.1	-0.2826219084407503	Iglesias, J., Vila, R., Canigueral, S., Bellakdhar, and II Idrissi, A. 1991. Analysis of the Essential Oil of <i>Thymus riatarum</i> . J. Ess. Oil Res. 3: 43-4.
<i>Thymus funkii</i>	Shoot		24.0	-0.20078804199300457	Vila, R., et al. 1995. Composition and study of the variability of the essential oil of <i>Thymus funkii</i> Cousson. Flav. & Fragr. J. 10(6): 379-383.
<i>Thymus longicaulis</i>	Shoot		1.7	-0.2771434906032444	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of <i>Thymus longicaulis</i> C. Presl subsp. <i>longicaulis</i> in the same Population. J. Ess. Oil Res. 5: 291-5.
<i>Thymus funkii</i>	Shoot		24.0	-0.20078804199300457	Vila, R., et al. 1995. Composition and study of the variability of the essential oil of <i>Thymus funkii</i> Cousson. Flav. & Fragr. J. 10(6): 379-383.
<i>Thymus longicaulis</i>	Shoot		0.0	-0.2829643095555945	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of <i>Thymus longicaulis</i> C. Presl subsp. <i>longicaulis</i> in the same Population. J. Ess. Oil Res. 5: 291-5.
<i>Thymus capitatus</i>	Shoot		0.0	-0.2829643095555945	Biondi, D., Cianci, P., Geraci, C. and Ruberto, G. 1993. Antimicrobial Activity and Chemical Composition of Essential Oils from Sicilian Aromatic Plants. Flav. & Fragr. J. 8: 331-7.
<i>Thymus longicaulis</i>	Shoot		0.0	-0.2829643095555945	Baser, K.H.C., Ozek, T., Kirimer, N. and Tumen, G. 1993. The Occurrence of Three Chemotypes of <i>Thymus longicaulis</i> C. Presl subsp. <i>longicaulis</i> in the same Population. J. Ess. Oil Res. 5: 291-5.
<i>Zingiber officinale</i>	Rhizome	2.0	115.0		--

Plant	Part	Low PPM	High PPM	StdDev	Reference
Zingiber officinale	Rhizome Essent. Oil		2300.0		--